

ANATOMY OF A q -GENERALIZATION OF THE LAGUERRE/HERMITE ORTHOGONAL POLYNOMIALS

CHUAN-TSUNG CHAN[†] AND HSIAO-FAN LIU^{††}

[†] *Department of Applied Physics, Tunghai University*

^{††} *Institute of Mathematics, Academia Sinica*

[†] *ctchan@go.thu.edu.tw*, ^{††} *liuhf@gate.sinica.edu.tw*

ABSTRACT. We study a q -generalization of the classical Laguerre/Hermite orthogonal polynomials. Explicit results include: the recursive coefficients, matrix elements of generators for the Heisenberg algebra, and the Hankel determinants. The power of quadratic relation is illustrated by comparing two ways of calculating recursive coefficients. Finally, we derive a q -deformed version of the Toda equations for both q -Laguerre/Hermite ensembles.

1. INTRODUCTION

One special property about the orthogonal polynomials is that the matrix representations of the Heisenberg algebra are usually very simple. For instance, the famous three-term recurrence relation among orthogonal polynomials

$$xp_n = a_{n+1}p_{n+1} + b_np_n + a_np_{n-1} \quad (1.1)$$

implies that the matrix form of the position operator, x , with respect to the orthogonal polynomial basis, is tri-diagonal for all kinds of weight functions. In addition, for many classical weight functions, one can show that both translation ($\frac{d}{dx}$) and dilation ($x\frac{d}{dx}$) operators are also tri-diagonal, such as the Laguerre and Hermite weight functions. In this paper, we shall extend these classical examples to a more general setting, which we call generalized q -Laguerre/Hermite ensembles, and give explicit results for the representation of the Heisenberg algebra. By *generalized*, we mean that, in addition to the exponential functions, we also multiply the *standard* Laguerre/Hermite weights by a polynomial prefactor. Or equivalently, we add an logarithmic term to the polynomial potential (see the definitions in Sec. 2).

Admittedly, the study of the Laguerre/Hermite ensembles might seem uninteresting for the apparent linearity features. However, from physical point of view, these two sets of orthogonal polynomials lie at the heart of fundamental physics. In particular, they are associated with the quantum systems of three dimensional hydrogen atom (associated Laguerre polynomials) and one dimensional simple harmonic oscillator (Hermite polynomials). Given the mysterious connection between these two systems, [9], it might be worthwhile to examine the physical relevance of the quadratic relation between these two ensembles (see Sec. 2). Furthermore, q -deformation are often considered as certain microscopic/quantum modifications

Date: December 30, 2016.

2010 Mathematics Subject Classification. 33D45, 39A45.

Key words and phrases. orthogonal polynomials, Laguerre polynomial, Hermite polynomial, Hankel determinants, Toda equation, matrix model.

of the existing physical models. Hence we would like to extend the classical scenarios to a possible q -generalizations.

To summarize, our new findings in this paper include:

- A systematic study of the generalized q -Laguerre and q -Hermite ensembles, including explicit solutions of the recursive coefficients, their orthogonal polynomial representations of the Heisenberg algebra, and their corresponding Hankel determinants (at $\kappa = 1$).
- We elucidate the power of the quadratic relation in comparing two ways of solving recursive coefficients for the q -Hermite ensemble.
- We find q -generalized Toda equations for the recursive coefficients of the q -Laguerre/Hermite ensembles.

2. THE QUADRATIC RELATIONS FOR THE LAGUERRE/HERMITE ORTHOGONAL POLYNOMIALS

2.1. Review of the quadratic relation among recursive coefficients for the orthogonal polynomials associated with the classical Laguerre/Hermite weights.

Given the classical Laguerre weight defined as

$$v^{(\alpha)}(x, \kappa) := x^\alpha \exp(-\kappa x), \quad 0 \leq \kappa, -1 < \alpha, 0 \leq x, \quad (2.1)$$

we can compute the orthonormal polynomials $p_n^{(\alpha)}(x, \kappa)$ as

$$\int_0^\infty p_m^{(\alpha)}(x, \kappa) p_n^{(\alpha)}(x, \kappa) v^{(\alpha)}(x, \kappa) dx = \delta_{mn} \quad (2.2)$$

through Gram-Schmidt process.

Similarly, from the classical Hermite weight,

$$\omega^{(\alpha)}(x, \kappa) := |x|^{2\alpha+1} \exp(-\kappa^2 x^2), \quad x \in \mathbb{R}, \quad (2.3)$$

we obtain associated orthonormal polynomials $P_n^{(\alpha)}(x, \kappa)$ as

$$\int_{-\infty}^\infty P_m^{(\alpha)}(x, \kappa) P_n^{(\alpha)}(x, \kappa) \omega^{(\alpha)}(x, \kappa) dx = \delta_{mn}. \quad (2.4)$$

The quadratic relation among these two sets of orthonormal polynomials is based on a simple connection between the Laguerre and Hermite weights. Namely,

$$\omega^{(\alpha)}(x, \kappa) = |x| v^{(\alpha)}(x^2, \kappa^2). \quad (2.5)$$

One immediate consequence of (2.5) is that, the orthonormal polynomials $P_n^{(\alpha)}(x, \kappa)$ can be expressed in terms of the orthogonal polynomials $p_n^{(\alpha)}(x, \kappa)$ as follows,

$$P_{2n}^{(\alpha)}(x, \kappa) = p_n^{(\alpha)}(x^2, \kappa^2), \quad P_{2n+1}^{(\alpha)}(x, \kappa) = x p_n^{(\alpha+1)}(x^2, \kappa^2). \quad (2.6)$$

Having established this fundamental relation, we can then derive many useful consequences. In this paper, we shall focus on quadratic relation for the recursive coefficients.

The set of orthonormal polynomials associated with any weight function can be viewed as a complete set of basis for the function space. Hence, it induces a natural realization of the Heisenberg algebra, $[\frac{d}{dx}, x] = 1$. In particular, the matrix elements of the position operator consist of the three-term recursive coefficients among orthonormal polynomials. In the case of the Laguerre weight, it is given as

$$xp_n^{(\alpha)}(x, \kappa) = a_{n+1}^{(\alpha)}(\kappa)p_{n+1}^{(\alpha)}(x, \kappa) + b_n^{(\alpha)}(\kappa)p_n^{(\alpha)}(x, \kappa) + a_n^{(\alpha)}(\kappa)p_{n-1}^{(\alpha)}(x, \kappa), \quad (2.7)$$

and in the case of the Hermite weight, we have

$$xP_n^{(\alpha)}(x, \kappa) = A_{n+1}^{(\alpha)}(\kappa)P_{n+1}^{(\alpha)}(x, \kappa) + A_n^{(\alpha)}(\kappa)P_{n-1}^{(\alpha)}(x, \kappa). \quad (2.8)$$

By computing $x^2P_n^{(\alpha)}(x, \kappa)$ in two ways, we obtain the quadratic relation among the two sets of recursive coefficients:

$$a_n^{(\alpha)}(\kappa^2) = A_{2n}^{(\alpha)}(\kappa)A_{2n-1}^{(\alpha)}(\kappa), \quad (2.9)$$

$$b_n^{(\alpha)}(\kappa^2) = \left(A_{2n+1}^{(\alpha)}(\kappa)\right)^2 + \left(A_{2n}^{(\alpha)}(\kappa)\right)^2. \quad (2.10)$$

2.2. On the quadratic relation between generalized q -Laguerre/Hermite ensembles.

In this paper, we take generalized q -Hermite and q -Laguerre ensembles as an illustrative example of the quadratic relation. In the simple case that $\kappa = 1$, we consider the q -Laguerre weight

$$v^{(\alpha)}(x; q) := |x|^\alpha (qx; q)_\infty = |x|^\alpha \prod_{k=0}^{\infty} (1 - q^{k+1}x), \quad (2.11)$$

and q -Hermite weight

$$\omega^{(\alpha)}(x; q) = |x|^{2\alpha+1} (q^2x^2; q^2)_\infty = |x|v^{(\alpha)}(x^2; q^2). \quad (2.12)$$

Given the orthonormal polynomials of the q -Laguerre ensemble $p_n^{(\alpha)}(x, q)$, we can express the orthonormal polynomials of the q -Hermite ensemble $P_n^{(\alpha)}(x, q)$ as follows:

$$P_{2n}^{(\alpha)}(x, q) = \sqrt{\frac{1+q}{2}} p_n^{(\alpha)}(x^2, q^2) \quad (\text{even, deg} = 2n), \quad (2.13)$$

$$P_{2n+1}^{(\alpha)}(x, q) = \sqrt{\frac{1+q}{2}} xp_n^{(\alpha+1)}(x^2, q^2) \quad (\text{odd, deg} = 2n+1). \quad (2.14)$$

One can easily check that $P_{2n}^{(\alpha)}(x, q)$ and $P_{2n+1}^{(\alpha)}(x, q)$ satisfy the orthonormal conditions, for instance,

$$\begin{aligned}
& \int_{-1}^1 P_{2m}^{(\alpha)}(x, q) P_{2n}^{(\alpha)}(x, q) \omega^{(\alpha)}(x, q) d_q x \\
&= 2(1-q) \sum_{k=0}^{\infty} P_{2m}^{(\alpha)}(q^k, q) P_{2n}^{(\alpha)}(q^k, q) \omega^{(\alpha)}(q^k, q) q^k \\
&= 2(1-q) \frac{1+q}{2} \sum_{k=0}^{\infty} p_m^{(\alpha)}(q^{2k}, q^2) p_n^{(\alpha)}(q^{2k}, q^2) q^k v^{(\alpha)}(q^{2k}, q^2) q^k \\
&= \int_0^1 p_m^{(\alpha)}(x, q^2) p_n^{(\alpha)}(x, q^2) v^{(\alpha)}(x, q^2) d_q x = \delta_{mn}. \tag{2.15}
\end{aligned}$$

$$\begin{aligned}
& \int_{-1}^1 P_{2m+1}^{(\alpha)}(x, q) P_{2n+1}^{(\alpha)}(x, q) \omega^{(\alpha)}(x, q) d_q x \\
&= 2(1-q) \sum_{k=0}^{\infty} P_{2m+1}^{(\alpha)}(q^k, q) P_{2n+1}^{(\alpha)}(q^k, q) \omega^{(\alpha)}(q^k, q) q^k \\
&= 2(1-q) \frac{1+q}{2} \sum_{k=0}^{\infty} q^k p_m^{(\alpha+1)}(q^{2k}, q^2) q^k p_n^{(\alpha+1)}(q^{2k}, q^2) q^k v^{(\alpha)}(q^{2k}, q^2) q^k \\
&= (1-q^2) \sum_{k=0}^{\infty} p_m^{(\alpha+1)}(q^{2k}, q^2) p_n^{(\alpha+1)}(q^{2k}, q^2) v^{(\alpha+1)}(q^{2k}, q^2) q^{2k} \\
&= \int_0^1 p_m^{(\alpha+1)}(x, q^2) p_n^{(\alpha+1)}(x, q^2) v^{(\alpha+1)}(x, q^2) d_q x = \delta_{mn}. \tag{2.16}
\end{aligned}$$

The $P_{2m}^{(\alpha)}-P_{2m+1}^{(\alpha)}$ orthogonality is trivial due to the even parity of q -Hermite weight.

By quadratic relation, we are mostly concerned with the correspondence between recursive coefficients associated with the q -Laguerre and q -Hermite ensembles. For example,

$$\begin{aligned}
x^2 P_{2m}^{(\alpha)}(x, q) &= x \left[A_{2m+1}^{(\alpha)}(q) P_{2m+1}^{(\alpha)}(x, q) + A_{2m}^{(\alpha)}(q) P_{2m-1}^{(\alpha)}(x, q) \right] \\
&= A_{2m+1}^{(\alpha)}(q) \left[A_{2m+2}^{(\alpha)}(q) P_{2m+2}^{(\alpha)}(x, q) + A_{2m+1}^{(\alpha)}(q) P_{2m-1}^{(\alpha)}(x, q) \right] \\
&\quad + A_{2m}^{(\alpha)}(q) \left[A_{2m}^{(\alpha)}(q) P_{2m}^{(\alpha)}(x, q) + A_{2m-1}^{(\alpha)}(q) P_{2m-2}^{(\alpha)}(x, q) \right] \\
&= \left[A_{2m+1}^{(\alpha)}(q) A_{2m+2}^{(\alpha)}(q) \right] P_{2m+2}^{(\alpha)}(x, q) + \left[\left(A_{2m+1}^{(\alpha)}(q) \right)^2 + \left(A_{2m}^{(\alpha)}(q) \right)^2 \right] P_{2m}^{(\alpha)}(x, q) \\
&\quad + [A_{2m-1}^{(\alpha)}(q) A_{2m}^{(\alpha)}(q)] P_{2m-2}^{(\alpha)}(x, q).
\end{aligned}$$

On the other hand, using the expression of (2.13), we have

$$\begin{aligned}
& x^2 P_{2m}^{(\alpha)}(x, q) \\
&= x^2 \sqrt{\frac{1+q}{2}} p_m^{(\alpha)}(x^2, q^2) \\
&= \sqrt{\frac{1+q}{2}} \left[a_{m+1}^{(\alpha)}(q^2) p_{m+1}^{(\alpha)}(x^2, q^2) + b_m^{(\alpha)}(q^2) p_m^{(\alpha)}(x^2, q^2) + a_m^{(\alpha)}(q^2) p_{m-1}^{(\alpha)}(x^2, q^2) \right] \\
&= a_{m+1}^{(\alpha)}(q^2) P_{2m+2}^{(\alpha)}(x, q) + b_m^{(\alpha)}(q^2) P_{2m}^{(\alpha)}(x, q) + a_m^{(\alpha)}(q^2) P_{2m-2}^{(\alpha)}(x, q).
\end{aligned}$$

By comparing the coefficients on both expressions, we get

$$a_m^{(\alpha)}(q^2) = A_{2m}^{(\alpha)}(q) A_{2m-1}^{(\alpha)}(q), \quad (2.17)$$

$$b_m^{(\alpha)}(q^2) = \left(A_{2m+1}^{(\alpha)}(q) \right)^2 + \left(A_{2m}^{(\alpha)}(q) \right)^2. \quad (2.18)$$

If we examine similar calculations for the odd q -Hermite orthonormal polynomials, we get

$$\begin{aligned}
x^2 P_{2m+1}^{(\alpha)} &= [A_{2m+3}^{(\alpha)} A_{2m+2}^{(\alpha)}] P_{2m+3}^{(\alpha)} \\
&+ \left[\left(A_{2m+2}^{(\alpha)} \right)^2 + \left(A_{2m+1}^{(\alpha)} \right)^2 \right] P_{2m+1}^{(\alpha)} + [A_{2m+1}^{(\alpha)} A_{2m}^{(\alpha)}] P_{2m-1}^{(\alpha)}.
\end{aligned}$$

Alternatively,

$$\begin{aligned}
& x^2 P_{2m+1}^{(\alpha)}(x, q) \\
&= x^2 \sqrt{\frac{1+q}{2}} x p_m^{(\alpha+1)}(x^2, q^2) \\
&= \sqrt{\frac{1+q}{2}} \left(a_{m+1}^{(\alpha+1)}(q^2) p_{m+1}^{(\alpha+1)}(x^2, q^2) + b_m^{(\alpha+1)}(q^2) p_m^{(\alpha+1)}(x^2, q^2) + a_m^{(\alpha+1)}(q^2) p_{m-1}^{(\alpha+1)}(x^2, q^2) \right) \\
&= a_{m+1}^{(\alpha+1)}(q^2) P_{2m+3}^{(\alpha)}(x, q) + b_m^{(\alpha+1)}(q^2) P_{2m+1}^{(\alpha)}(x, q) + a_m^{(\alpha+1)}(q^2) P_{2m-1}^{(\alpha)}(x, q).
\end{aligned}$$

Thus,

$$a_m^{(\alpha+1)}(q^2) = A_{2m+1}^{(\alpha)}(q) A_{2m}^{(\alpha)}(q), \quad (2.19)$$

$$b_m^{(\alpha+1)}(q^2) = \left(A_{2m+2}^{(\alpha)}(q) \right)^2 + \left(A_{2m+1}^{(\alpha)}(q) \right)^2. \quad (2.20)$$

Eliminate the recursive coefficients of the q -Laguerre orthonormal polynomials, $a_m^{(\alpha)}, b_m^{(\alpha)}$, in both sets of the equation, we obtain

$$A_{2m}^{(\alpha)}(q) A_{2m-2}^{(\alpha)}(q) = A_{2m+1}^{(\alpha-1)}(q) A_{2m}^{(\alpha-1)}(q), \quad (2.21)$$

and

$$\left(A_{2m+1}^{(\alpha)}(q)\right)^2 + \left(A_{2m}^{(\alpha)}(q)\right)^2 = \left(A_{2m+2}^{(\alpha-1)}(q)\right)^2 + \left(A_{2m+1}^{(\alpha-1)}(q)\right)^2. \quad (2.22)$$

We shall see how these quadratic relation will help us derive the explicit form of the recursive coefficients $A_n^{(\alpha)}(q)$ from the (easier) solutions of $a_n^{(\alpha)}(q^2), b_n^{(\alpha)}(q^2)$ in Sec. 3.

3. EXPLICIT SOLUTIONS OF THE FOURIER AND RECURSIVE COEFFICIENTS FOR THE q -LAGUERRE/HERMITE ENSEMBLES

3.1. Solution of the recursive coefficients for the generalized q -Laguerre ensemble (at $\kappa = 1$).

To compute the recursive coefficients, $a_n^{(\alpha)}(q), b_n^{(\alpha)}(q)$, we first study the Fourier coefficients of the action of dilation on the orthonormal q -Laguerre polynomials:

$$x\mathcal{D}_q^x p_n^{(\alpha)}(x) = \sum_{j=0}^n p_j^{(\alpha)}(x) c_{jn}^{(\alpha)}, \quad (3.1)$$

where

$$\begin{aligned} c_{jn}^{(\alpha)} &= \int_{-1}^1 p_j^{(\alpha)}(x) [x\mathcal{D}_q^x p_n^{(\alpha)}(x)] v^{(\alpha)}(x) d_q x \\ &= \frac{-1}{1-q} \int_{-1}^1 p_j^{(\alpha)}(x) p_n^{(\alpha)}(qx) v^{(\alpha)}(x) d_q x \\ &\quad + \frac{1}{1-q} \int_{-1}^1 p_j^{(\alpha)}(x) p_n^{(\alpha)}(x) v^{(\alpha)}(x) d_q x. \end{aligned} \quad (3.2)$$

The Pearson relation for the generalized q -Laguerre weight,

$$v^{(\alpha)}\left(\frac{x}{q}\right) = q^{-\alpha}(1-x)v^{(\alpha)}(x), \quad (3.3)$$

we get

$$\begin{aligned} c_{jn}^{(\alpha)} &= -\frac{q^{-(\alpha+1)}}{1-q} \int_{-1}^1 p_j^{(\alpha)}\left(\frac{x}{q}\right) p_n^{(\alpha)}(x) v^{(\alpha)}(x) d_q x \\ &\quad + \frac{q^{-(\alpha+1)}}{1-q} \int_{-1}^1 p_j^{(\alpha)}\left(\frac{x}{q}\right) p_n^{(\alpha)}(x) x v^{(\alpha)}(x) d_q x \\ &\quad + \frac{1}{1-q} \delta_{jn}. \end{aligned} \quad (3.4)$$

From this expression, it is clear that $c_{jn}^{(\alpha)} = 0$ if $j < n - 1$. Thus, we have

$$x\mathcal{D}_q^x p_n^{(\alpha)}(x) = p_n^{(\alpha)}(x)c_{nn}^{(\alpha)} + p_{n-1}^{(\alpha)}(x)c_{n-1,n}^{(\alpha)}. \quad (3.5)$$

From the coefficients of the orthonormal q -Laguerre polynomials:

$$p_n^{(\alpha)}(x) = \gamma_n^{(\alpha)}x^n + \delta_n^{(\alpha)}x^{n-1} + \cdots, \quad \eta_n^{(\alpha)} := \frac{\delta_n^{(\alpha)}}{\gamma_n^{(\alpha)}} \quad (3.6)$$

we deduce the decomposition of the rescaled orthonormal polynomials,

$$\begin{aligned} p_n^{(\alpha)}\left(\frac{x}{q}\right) &= q^{-n}p_n^{(\alpha)}(x) + \frac{\eta_n^{(\alpha)}}{a_n^{(\alpha)}}q^{-n}(q-1)p_{n-1}^{(\alpha)}(x) + \cdots, \\ p_n^{(\alpha)}(qx) &= q^n p_n^{(\alpha)}(x) + \frac{\eta_n^{(\alpha)}}{a_n^{(\alpha)}}q^{n-1}(1-q)p_{n-1}^{(\alpha)}(x) + \cdots. \end{aligned} \quad (3.7)$$

Substituting these results back to Eq.(3.2), we obtain

$$c_{nn}^{(\alpha)} = \frac{1-q^n}{1-q} \text{ (independent of } \alpha), \quad (3.8)$$

$$c_{n-1,n}^{(\alpha)} = -\frac{\eta_n^{(\alpha)}}{a_n^{(\alpha)}}q^{n-1}. \quad (3.9)$$

On the other hand, we can relate the Fourier coefficients, $c_{nn}^{(\alpha)}$ and $c_{n-1,n}^{(\alpha)}$ to the recursive coefficient $a_n^{(\alpha)}$ using the second integral expression together with

$$xp_n^{(\alpha)}\left(\frac{x}{q}\right) = q^{-n}a_{n+1}^{(\alpha)}p_{n+1}^{(\alpha)}(x) + (\eta_n^{(\alpha)}q^{-(n+1)} - \eta_{n+1}^{(\alpha)}q^{-n})p_n^{(\alpha)}(x) + \cdots. \quad (3.10)$$

The results are:

$$c_{nn}^{(\alpha)} = -\frac{q^{-(\alpha+n+1)}}{1-q} + \frac{q^{-(\alpha+1)}}{1-q}(\eta_n^{(\alpha)}q^{-(n-1)} - \eta_{n+1}^{(\alpha)}q^{-n}) + \frac{1}{1-q}, \quad (3.11)$$

$$c_{n-1,n}^{(\alpha)} = \frac{q^{-(\alpha+n)}}{1-q}a_n^{(\alpha)}. \quad (3.12)$$

By comparing two expressions for $c_{nn}^{(\alpha)}$, Eqs.(3.9) and (3.12), we solve

$$\eta_n^{(\alpha)} = \frac{(q^{\alpha+n} - 1)(1 - q^n)}{1 - q}. \quad (3.13)$$

From the expressions for $c_{n-1,n}^{(\alpha)}$, Eqs.(3.9) and (3.12), we solve

$$(a_n^{(\alpha)})^2 = q^{\alpha+2n-1}(1 - q^n)(1 - q^{\alpha+n}) \geq 0. \quad (3.14)$$

Finally, the middle recursive coefficient $b_n^{(\alpha)}$ can be solved using the formula in the Appendix:

$$b_n^{(\alpha)} = \eta_n^{(\alpha)} - \eta_{n+1}^{(\alpha)} = -q^{\alpha+2n}(1+q) + q^{\alpha+n} + q^n. \quad (3.15)$$

3.2. Solution of the recursive coefficients for the generalized q -Hermite ensemble (at $\kappa = 1$).

In this section, we calculate the recursive coefficients for the orthonormal polynomials associated with generalized q -Hermite ensembles, $A_n^{(\alpha)}$.

In order to derive a master equation for the recursive coefficients $A_n^{(\alpha)}$, we first study the Fourier coefficients of the action of dilation on the orthonormal q -Hermite polynomials

$$x\mathcal{D}_q^x P_n^{(\alpha)}(x) = \sum_{j=0}^n P_j^{(\alpha)}(x) C_{jn}^{(\alpha)}, \quad (n-j \text{ must be even}) \quad (3.16)$$

where

$$C_{jn}^{(\alpha)} = \int_{-1}^1 P_j^{(\alpha)}(x) [x\mathcal{D}_q^x P_n^{(\alpha)}(x)] \omega^{(\alpha)}(x) d_q x \quad (3.17)$$

Substituting the definition of the q -derivative $\mathcal{D}_q^x P_n^{(\alpha)}$ and recalling the definition of the q -integration, we get

$$\begin{aligned} C_{jn}^{(\alpha)} &= -2 \sum_{l=0}^{\infty} P_j^{(\alpha)}(q^l) P_n^{(\alpha)}(q^{l+1}) \omega^{(\alpha)}(q^l) q^l \\ &+ 2 \sum_{l=0}^{\infty} P_j^{(\alpha)}(q^l) P_n^{(\alpha)}(q^l) \omega^{(\alpha)}(q^l) q^l \\ &= \frac{-1}{1-q} \int_{-1}^1 P_j^{(\alpha)}(x) P_n^{(\alpha)}(qx) \omega^{(\alpha)}(x) d_q x \\ &+ \frac{1}{1-q} \int_{-1}^1 P_j^{(\alpha)}(x) P_n^{(\alpha)}(x) \omega^{(\alpha)}(x) d_q x. \end{aligned} \quad (3.18)$$

Upon shifting the index κ and using the Pearson relation for the generalized q -Hermite weight

$$\omega^{(\alpha)}\left(\frac{x}{q}\right) = q^{-(2\alpha+1)}(1-x^2)\omega^{(\alpha)}(x), \quad (3.19)$$

we get

$$\begin{aligned}
C_{jn}^{(\alpha)} &= -\frac{q^{-(2\alpha+2)}}{1-q} \int_{-1}^1 P_j^{(\alpha)}\left(\frac{x}{q}\right) P_n^{(\alpha)}(x) \omega^{(\alpha)}(x) d_q x \\
&+ \frac{q^{-(2\alpha+2)}}{1-q} \int_{-1}^1 P_j^{(\alpha)}\left(\frac{x}{q}\right) P_n^{(\alpha)}(x) x^2 \omega^{(\alpha)}(x) d_q x \\
&+ \frac{1}{1-q} \delta_{jn}.
\end{aligned} \tag{3.20}$$

From this expression, it is clear that $C_{jn}^{(\alpha)} = 0$ if $j < n - 2$. Thus, we have

$$x \mathcal{D}_q^x P_n^{(\alpha)}(x) = P_n^{(\alpha)}(x) C_{nn}^{(\alpha)} + P_{n-2}^{(\alpha)}(x) C_{n-2,n}^{(\alpha)}. \tag{3.21}$$

To facilitate the computations, we need to invoke the structure of the orthonormal coefficients. Let

$$P_n^{(\alpha)}(x) = \gamma_n^{(\alpha)} x^n + \delta_n^{(\alpha)} x^{n-2} + \dots, \quad \zeta_n^{(\alpha)} := \frac{\delta_n^{(\alpha)}}{\gamma_n^{(\alpha)}},$$

we can decompose the rescaled orthonormal polynomials $P_n^{(\alpha)}\left(\frac{x}{q}\right)$, $P_n^{(\alpha)}(qx)$ as follows:

$$\begin{aligned}
P_n^{(\alpha)}\left(\frac{x}{q}\right) &= \gamma_n^{(\alpha)} q^{-n} x^n + \delta_n^{(\alpha)} q^{-n+2} x^{n-2} + \dots \\
&= q^{-n} (\gamma_n^{(\alpha)} x^n + \delta_n^{(\alpha)} x^{n-2} + \dots) - q^{-n} \delta_n^{(\alpha)} x^{n-2} + \delta_n^{(\alpha)} q^{-n+2} x^{n-2} + \dots \\
&= q^{-n} P_n^{(\alpha)}(x) - \frac{q^{-n}(1-q^2)\zeta_n^{(\alpha)}}{A_{n-1}^{(\alpha)} A_n^{(\alpha)}} P_{n-2}^{(\alpha)} + \dots,
\end{aligned} \tag{3.22}$$

$$\begin{aligned}
P_n^{(\alpha)}(qx) &= \gamma_n^{(\alpha)} q^n x^n + \delta_n^{(\alpha)} q^{n-2} x^{n-2} + \dots \\
&= q^n (\gamma_n^{(\alpha)} x^n + \delta_n^{(\alpha)} x^{n-2} + \dots) - q^n \delta_n^{(\alpha)} x^{n-2} + \delta_n^{(\alpha)} q^{n-2} x^{n-2} + \dots \\
&= q^n P_n^{(\alpha)}(x) + \frac{q^{n-2}(1-q^2)\zeta_n^{(\alpha)}}{A_{n-1}^{(\alpha)} A_n^{(\alpha)}} P_{n-2}^{(\alpha)} + \dots.
\end{aligned} \tag{3.23}$$

Substituting these results back to (3.18), we obtain

$$C_{nn}^{(\alpha)} = \frac{1-q^n}{1-q} \text{ (independent of } \alpha), \tag{3.24}$$

$$C_{n-2,n}^{(\alpha)} = \frac{-(1+q)q^{n-2}}{A_{n-1}^{(\alpha)} A_n^{(\alpha)}} \zeta_n^{(\alpha)}. \tag{3.25}$$

Note that

$$\zeta_n^{(\alpha)} := \frac{\delta_n^{(\alpha)}}{\gamma_n^{(\alpha)}} = - \sum_{l=0}^{n-1} \left(A_l^{(\alpha)} \right)^2, \quad (3.26)$$

and

$$C_{n+1,n+1}^{(\alpha)} - C_{nn}^{(\alpha)} = q^n. \quad (3.27)$$

Having derived the Fourier coefficients $C_{nn}^{(\alpha)}$ and $C_{n-2,n}^{(\alpha)}$ for $x\mathcal{D}_q^x[P_n^{(\alpha)}(x)]$, we can use Eq.(3.20) to derive a master equation for the recursive coefficients $A_n^{(\alpha)}$ defined in Eq.(2.8) with $\kappa = 1$,

$$xP_n^{(\alpha)}(x) = A_{n+1}^{(\alpha)}P_{n+1}^{(\alpha)}(x) + A_n^{(\alpha)}P_{n-1}^{(\alpha)}(x). \quad (3.28)$$

Namely,

$$xP_n^{(\alpha)}\left(\frac{x}{q}\right) = P_{n+1}^{(\alpha)}(x)[q^{-n}A_{n+1}^{(\alpha)}] + P_{n-1}^{(\alpha)}(x)\left[q^{-n}A_n^{(\alpha)} - \frac{q^{-n}\zeta_n^{(\alpha)}(1-q^2)}{A_n^{(\alpha)}}\right] + \cdots, \quad (3.29)$$

Eq.(3.20) implies

$$\begin{aligned} 1 - q^n &= (1 - q)C_{nn}^{(\alpha)} \\ &= (1 - q^{-(2\alpha+2)-n}) + q^{-(2\alpha+2)-n} \left[\left(A_{n+1}^{(\alpha)} \right)^2 + \left(A_n^{(\alpha)} \right)^2 - \zeta_n^{(\alpha)}(1 - q^2) \right]. \end{aligned} \quad (3.30)$$

By taking a first difference on both sides of the equation, $(1 - q)(C_{n+1,n+1}^{(\alpha)} - C_{nn}^{(\alpha)})$, and define a shifted difference

$$B_k^{(\alpha)} := \left(A_k^{(\alpha)} \right)^2 - q \left(A_{k-1}^{(\alpha)} \right)^2 \Leftrightarrow \left(A_n^{(\alpha)} \right)^2 = \sum_{k=1}^n q^{n-k} B_k^{(\alpha)}, \quad (3.31)$$

with $A_0^{(\alpha)} = 0, B_1^{(\alpha)} = \left(A_1^{(\alpha)} \right)^2$, we get

$$(1 - q)q^{2n+2\alpha+3} = (q - 1) + B_{n+2}^{(\alpha)} + B_{n+1}^{(\alpha)} + (1 - q^2) \sum_{k=1}^n B_k^{(\alpha)}. \quad (3.32)$$

By taking another difference $(1 - q)(q^{2n+2\alpha+3} - q^{2n+2\alpha+1})$, and defining another shifted difference $C_n^{(\alpha)} := B_{n+1}^{(\alpha)} - qB_n^{(\alpha)}$. That is,

$$B_n^{(\alpha)} = q^{n-1}B_1^{(\alpha)} + \sum_{k=1}^{n-1} q^{n-1-k}C_k^{(\alpha)}, \quad (3.33)$$

we get

$$C_{n+1}^{(\alpha)} = -qC_n^{(\alpha)} - (1 - q)^2(1 + q)q^{2n+2\alpha+1}. \quad (3.34)$$

The explicit form of $C_n^{(\alpha)}$ can be solved by iteration, and is given by

$$C_n^{(\alpha)} = (-q)^{n-1} C_1^{(\alpha)} - (1-q)^2 (-1)^n q^{n+2\alpha+1} [1 - (-q)^{n-1}], \quad (3.35)$$

where $C_1^{(\alpha)} = B_2^{(\alpha)} - qB_1^{(\alpha)} = \left(A_2^{(\alpha)}\right)^2 - 2q \left(A_1^{(\alpha)}\right)^2$.

Substituting the explicit form of $C_n^{(\alpha)}$, and performing the summation, Eq.(3.33), we get for $m \geq 1$,

$$B_{2m}^{(\alpha)} = -q^{2m-1} \left(A_1^{(\alpha)}\right)^2 + q^{2m-2} \left(A_2^{(\alpha)}\right)^2 - q^{2m+\rho} (1-q)(1-q^{2m-2}), \quad (3.36)$$

$$B_{2m-1}^{(\alpha)} = q^{2m-2} \left(A_1^{(\alpha)}\right)^2 - q^{2m+\rho-2} (1-q)(1-q^{2m-2}). \quad (3.37)$$

Substituting the explicit forms of $B_n^{(\alpha)}$, and performing the summation, (3.31), we get

$$\left(A_{2m}^{(\alpha)}\right)^2 = mq^{2m-2} \left(A_2^{(\alpha)}\right)^2 + q^{2m+\rho-1} [(1-q^{2m}) - m(1-q^2)], \quad (3.38)$$

$$\begin{aligned} \left(A_{2m-1}^{(\alpha)}\right)^2 &= q^{2m-2} \left(A_1^{(\alpha)}\right)^2 + (m-1)q^{2m-3} \left(A_2^{(\alpha)}\right)^2 \\ &+ \frac{q^{2m+\rho-1}}{1+q} [(1-q^{2m-2}) - (m-1)(1-q)]. \end{aligned} \quad (3.39)$$

3.3. The power of the quadratic relation.

To show the power of the quadratic relation, we first use the solutions of the recursive coefficients of the q -Laguerre orthonormal polynomials $a_n^{(\alpha)}, b_n^{(\alpha)}$, Eqs.(3.14), (3.15) to deduce the recursive coefficients of the q -Hermite polynomials. First of all, we have learned that, using Eqs. (2.17), (2.18),

$$\begin{aligned} \left(A_{2n}^{(\alpha)}(q)\right)^2 \left(A_{2n+1}^{(\alpha)}(q)\right)^2 &= \left(a_n^{(\alpha+1)}(q^2)\right)^2 \\ &= q^{2\alpha+4n} (1-q^{2n}) (1-q^{2\alpha+2n+2}), \end{aligned}$$

and

$$\begin{aligned} \left(A_{2n}^{(\alpha)}(q)\right)^2 + \left(A_{2n+1}^{(\alpha)}(q)\right)^2 &= b_n^\alpha(q^2) \\ &= -q^{2\alpha+4n} (1+q^{2n}) + q^{2n} (q^{2\alpha} + 1). \end{aligned}$$

This implies that $\left(A_{2n}^{(\alpha)}\right)^2$ and $\left(A_{2n+1}^{(\alpha)}\right)^2$ can be solved from the quadratic equation,

$$z^2 - b_n^\alpha(q^2)z + \left(a_n^{(\alpha+1)}(q^2)\right)^2 = 0, \quad (3.40)$$

and two roots are

$$z_{\pm}^{(\alpha)}(q) = \frac{b_n^{(\alpha)}(q^2) \pm \sqrt{\left(b_n^{(\alpha)}(q^2)\right)^2 - 4\left(a_n^{(\alpha+1)}(q^2)\right)^2}}{2}. \quad (3.41)$$

Substituting the solutions for $a_n^{(\alpha+1)}(q^2)$ and $b_n^{(\alpha)}(q^2)$, Eqs.(3.14),(3.15), we get

$$\begin{aligned} z_{\pm}^{(\alpha)}(q) &= \frac{-q^{2\alpha+4n}(1+q^2) + q^2n(1+q^{2\alpha} \pm [q^{2\alpha+4n}(1-q^2) + q^{2n}(1-q^{2\alpha})])}{2} \\ &= \begin{cases} -q^{2\alpha+4n+2} + q^{2n} \\ -q^{2\alpha+4n} + q^{2\alpha+2n}. \end{cases} \end{aligned}$$

By taking $\alpha \rightarrow -\frac{1}{2}$,

$$z_{\pm}^{(-\frac{1}{2})}(q) \rightarrow \begin{cases} -q^{4n+1} + q^{2n} \rightarrow A_{2n+1}^{(-\frac{1}{2})} \\ -q^{4n-1} + q^{2n-1} \rightarrow A_{2n}^{(-\frac{1}{2})}. \end{cases} \quad (3.42)$$

We deduce that

$$\left(A_{2n}^{(\alpha)}\right)^2 = q^{2n+2\alpha}(1-q^{2n}) \Rightarrow \left(A_2^{(\alpha)}\right)^2 = q^{2\alpha+2}(1-q^2) \quad (3.43)$$

$$\left(A_{2n+1}^{(\alpha)}\right)^2 = q^{2n}(1-q^{2\alpha+2n+2}) \Rightarrow \left(A_1^{(\alpha)}\right)^2 = 1 - q^{2\alpha+2}. \quad (3.44)$$

From this second approach, it is clear that not only we can easily derive the recursive coefficients of the generalized q -Hermite orthonormal polynomials, but also we get the *initial* values $A_1^{(\alpha)}, A_2^{(\alpha)}$ for free. One can now check by substituting the *initial* values $A_1^{(\alpha)}, A_2^{(\alpha)}$ into Eqs.(3.38), (3.39), which are solutions from (laborious) successive reduction, that we obtain consistent results as derived from the quadratic relation.

4. HANKEL DETERMINANTS AND ITS q -GENERALIZATION

4.1. Hankel determinant for the q -Laguerre ensemble.

In this section, we shall compute the Hankel determinant associated with the generalized q -Laguerre ensemble, which is defined as the determinant of the moments associated with the weight function.

$$\mu_k^{(\alpha)} := \int_0^1 x^k v^{(\alpha)}(x, q) d_q x, \quad (4.1)$$

$$\Delta_n^{(\alpha)} := \begin{vmatrix} \mu_0 & \mu_1 & \cdots & \mu_{n-1} \\ \mu_1 & \mu_2 & \cdots & \mu_n \\ \vdots & \vdots & & \vdots \\ \mu_{n-1} & \mu_n & \cdots & \mu_{2n-2} \end{vmatrix}, \quad \Delta_0^{(\alpha)} := 1. \quad (4.2)$$

$$\tilde{\Delta}_n^{(\alpha)} := \begin{vmatrix} \mu_0 & \mu_1 & \cdots & \mu_{n-2} & \mu_n \\ \mu_1 & \mu_2 & \cdots & \mu_{n-1} & \mu_{n+1} \\ \vdots & \vdots & & \vdots & \vdots \\ \mu_{n-1} & \mu_n & \cdots & \mu_{2n-3} & \mu_{2n-1} \end{vmatrix}, \quad \tilde{\Delta}_0^{(\alpha)} := 1. \quad (4.3)$$

In addition, the recursive coefficients for the orthonormal polynomials can be expressed in terms of the Hankel determinants as

$$(a_n^{(\alpha)})^2 = \frac{\Delta_{n+1}^{(\alpha)} \Delta_{n-1}^{(\alpha)}}{(\Delta_n^{(\alpha)})^2}, \quad b_n^{(\alpha)} = \frac{\tilde{\Delta}_{n+1}^{(\alpha)}}{\Delta_{n+1}^{(\alpha)}} - \frac{\tilde{\Delta}_n^{(\alpha)}}{\Delta_n^{(\alpha)}}. \quad (4.4)$$

Defining the ratio among the Hankel determinants of adjacent rank,

$$R_n^{(\alpha)} := \frac{\Delta_n^{(\alpha)}}{\Delta_{n-1}^{(\alpha)}}, \quad R_1^{(\alpha)} = \frac{\Delta_1^{(\alpha)}}{\Delta_0^{(\alpha)}} = \mu_0^{(\alpha)} = \frac{1}{(\gamma_0^{(\alpha)})^2}, \quad (4.5)$$

we see that

$$(a_n^{(\alpha)})^2 = \frac{R_{n+1}^{(\alpha)}}{R_n^{(\alpha)}}, \quad \text{and} \quad R_n^{(\alpha)} = R_1^{(\alpha)} \prod_{k=1}^{n-1} (a_k^{(\alpha)})^2. \quad (4.6)$$

Now we can substitute the solutions of the recursive coefficients, Eq.(3.14), for $(a_k^{(\alpha)})^2$ and we get

$$\begin{aligned} R_n^{(\alpha)} &= R_1^{(\alpha)} q^{\sum_{k=1}^{n-1} (\alpha+2k-1)} \left(\prod_{k=1}^{n-1} (1-q^k)(1-q^{\alpha+k}) \right) \\ &= R_1^{(\alpha)} q^{(n-1)(n+\alpha-1)} \left(\prod_{l=1}^{n-1} (1-q^l)(1-q^{\alpha+l}) \right). \end{aligned} \quad (4.7)$$

Next, we compute the Hankel determinant

$$\begin{aligned} \Delta_n^{(\alpha)} &= \Delta_1^{(\alpha)} \prod_{k=2}^n R_k^{(\alpha)} \\ &= \Delta_1^{(\alpha)} \left(R_1^{(\alpha)} \right)^{n-1} q^{\sum_{k=2}^n (k-1)(k+\alpha-1)} \prod_{k=1}^{n-1} \prod_{l=1}^k ((1-q^l)(1-q^{l+\alpha})) \\ &= \left(\Delta_1^{(\alpha)} \right)^n q^{\frac{n(n-1)(2n-1+3\alpha)}{6}} \prod_{k=1}^{n-1} (1-q^k)^{n-k} (1-q^{k+\alpha})^{n-k}. \end{aligned} \quad (4.8)$$

4.2. Hankel determinant for the q -Hermite ensemble.

Following the same definitions, we compute the Hankel determinant associated with the q -Hermite ensemble. We recall the solutions of the recursive coefficients, Eqs.(3.43),(3.44),

$$\begin{aligned} \left(A_{2k}^{(\alpha)}(q)\right)^2 &= q^{2(k+\alpha)}(1 - q^{2k}), \\ \left(A_{2k-1}^{(\alpha)}(q)\right)^2 &= q^{2(k-1)}(1 - q^{2(k+\alpha)}), \end{aligned}$$

and define the ratios among adjacent Hankel determinants, $R_n^{(\alpha)} := \frac{\Delta_n^{(\alpha)}}{\Delta_{n-1}^{(\alpha)}}$, we get

$$\begin{aligned} R_{2m}^{(\alpha)} &= R_1^{(\alpha)} \prod_{l=1}^{2m-1} \left(A_l^{(\alpha)}\right)^2 \\ &= R_1^{(\alpha)} \left\{ \prod_{k=1}^{m-1} \left(A_{2k}^{(\alpha)}\right)^2 \right\} \left\{ \prod_{l=1}^m \left(A_{2l-1}^{(\alpha)}\right)^2 \right\} \\ &= R_1^{(\alpha)} q^{2 \sum_{k=1}^{m-1} (k+\alpha)} \left(\prod_{k=1}^{m-1} (1 - q^{2k}) \right) q^{2 \sum_{l=1}^m (l-1)} \left(\prod_{l=1}^m (1 - q^{2(l+\alpha)}) \right) \\ &= R_1^{(\alpha)} q^{2(m+\alpha)(m-1)} \prod_{k=1}^{m-1} (1 - q^{2k}) \prod_{l=1}^m (1 - q^{2(l+\alpha)}). \end{aligned} \tag{4.9}$$

Similarly,

$$\begin{aligned} R_{2m-1}^{(\alpha)} &= R_1^{(\alpha)} \prod_{l=1}^{2m-2} \left(A_l^{(\alpha)}\right)^2 \\ &= R_1^{(\alpha)} \left\{ \prod_{k=1}^{m-1} \left(A_{2k}^{(\alpha)}\right)^2 \right\} \left\{ \prod_{l=1}^{m-1} \left(A_{2l-1}^{(\alpha)}\right)^2 \right\} \\ &= R_1^{(\alpha)} q^{2(\alpha+m)(m-1)} \left(\prod_{k=1}^{m-1} (1 - q^{2k}) \right) q^{(m-2)(m-1)} \left(\prod_{l=1}^{m-1} (1 - q^{2(l+\alpha)}) \right) \\ &= R_1^{(\alpha)} q^{2(m-1)(m+\alpha-1)} \prod_{k=1}^{m-1} (1 - q^{2k}) (1 - q^{2(k+\alpha)}). \end{aligned} \tag{4.10}$$

From the results of Eqs.(4.9), (4.10), we can then derive explicit expressions for the Hankel determinants as a finite product. For the even case,

$$\begin{aligned}
\Delta_{2j}^{(\alpha)} &= \Delta_1^{(\alpha)} \prod_{k=2}^{2j} R_k^{(\alpha)} \\
&= \Delta_1^{(\alpha)} \left(\prod_{k=1}^j R_{2k}^{(\alpha)} \right) \left(\prod_{l=1}^{j-1} R_{2l+1}^{(\alpha)} \right) \\
&= \left(\Delta_1^{(\alpha)} \right)^{2j} q^{\frac{j(j-1)(4j+6\alpha+1)}{3}} \left(\prod_{k=1}^{j-1} (1 - q^{2k})^{2j-2k} \right) \left(\prod_{l=1}^j (1 - q^{2l+2\alpha})^{2j-2l+1} \right). \quad (4.11)
\end{aligned}$$

The odd case can be easily computed

$$\begin{aligned}
\Delta_{2j-1}^{(\alpha)} &= \frac{\Delta_{2j}^{(\alpha)}}{R_{2j}^{(\alpha)}} \\
&= \left(\Delta_1^{(\alpha)} \right)^{2j-1} q^{\frac{j-1}{3} [4j^2 + (6\alpha-5)j - 6\alpha]} \left(\prod_{k=1}^{j-1} (1 - q^{2k})^{2j-2k-1} (1 - q^{2k+2\alpha})^{2j-2k} \right).
\end{aligned}$$

As a consistent check, we specialize our results to $\alpha = 0$ and $\alpha = -\frac{1}{2}$.

$\alpha = 0$:

$$R_{2j}^{(0)} = R_1^{(0)} q^{2j(j-1)} (1 - q^{2j}) \prod_{k=1}^{j-1} (1 - q^{2k})^2 \quad (4.12)$$

$$R_{2j-1}^{(0)} = R_1^{(0)} q^{2(j-1)^2} \prod_{k=1}^{j-1} (1 - q^{2k})^2 \quad (4.13)$$

$$\Delta_{2j}^{(0)} = \left(\Delta_1^{(0)} \right)^{2j} q^{\frac{j(j-1)(4j+1)}{3}} \prod_{k=1}^{j-1} ((1 - q^{2k})^{4j-4k+1})^{(1-q^{2j})} \quad (4.14)$$

$$\Delta_{2j-1}^{(0)} = \left(\Delta_1^{(0)} \right)^{2j-1} q^{\frac{j(j-1)}{3} (4j-5)} \prod_{k=1}^{j-1} (1 - q^{2k})^{4j-4k-1} \quad (4.15)$$

$$\alpha = -\frac{1}{2} :$$

$$R_{2j}^{(-\frac{1}{2})} = R_1^{(-\frac{1}{2})} q^{(2j-1)(j-1)} \prod_{k=1}^{2j-1} (1 - q^k) \quad (4.16)$$

$$R_{2j-1}^{(-\frac{1}{2})} = R_1^{(-\frac{1}{2})} q^{(2j-3)(j-1)} \prod_{k=1}^{2j-2} (1 - q^k) \quad (4.17)$$

$$\Delta_{2j}^{(-\frac{1}{2})} = \left(R_1^{(-\frac{1}{2})}\right)^{2j} q^{\frac{2j(j-1)(2j-1)}{3}} \prod_{k=1}^{2j-1} ((1 - q^k)^{2j-k}) \quad (4.18)$$

$$\Delta_{2j-1}^{(-\frac{1}{2})} = \left(R_1^{(-\frac{1}{2})}\right)^{2j-1} q^{\frac{(j-1)(2j-1)(2j-3)}{3}} \prod_{k=1}^{2j-2} (1 - q^k)^{2j-k-1} \quad (4.19)$$

5. q -GENERALIZATION OF THE TODA EQUATIONS FROM κ -DEFORMATION OF THE q -LAGUERRE/HERMITE ENSEMBLES

5.1. q -Difference equations for the recursive coefficients of the q -Laguerre polynomials.

In this section, we first compute the Fourier expansion (w.r.t κ variable) of the q -derivative on the q -Laguerre orthonormal polynomials. The Fourier coefficients are defined by

$$\mathcal{D}_q^\kappa p_n^{(\alpha)}(x, \kappa) = \sum_{j=0}^n p_j^{(\alpha)}(x, \kappa) \xi_{jn}^{(\alpha)}(\kappa). \quad (5.1)$$

Recalling the definition of the q -derivative, (A.2), we can also transform this expansion formula as a q -shifting relation ($\lambda = (1 - q)\kappa$):

$$p_n^{(\alpha)}(x, q\kappa) = p_n^{(\alpha)}(x, \kappa) [1 - \lambda \xi_{nn}^{(\alpha)}(\kappa)] - \lambda \sum_{j=0}^{n-1} p_j^{(\alpha)}(x, \kappa) \xi_{jn}^{(\alpha)}(\kappa). \quad (5.2)$$

By taking q -derivative w.r.t κ variable on the orthonormal condition, and recalling the q -Leibniz rule, (A.6), we can derive a master equation among these Fourier coefficients.

$$\mathcal{D}_q^\kappa \left[\int_0^1 p_m^{(\alpha)}(x, \kappa) p_n^{(\alpha)}(x, \kappa) v^{(\alpha)}(x, \kappa) d_q x \right] = 0. \quad (5.3)$$

This implies for $m \leq n$,

$$(1 - \lambda \xi_{mm}^{(\alpha)}) \xi_{mn}^{(\alpha)} - \lambda \sum_{j=0}^{m-1} \xi_{jm}^{(\alpha)} \xi_{jn}^{(\alpha)} + \delta_{mn} \xi_{nn}^{(\alpha)} = \delta_{m,n-1} \left[\frac{q}{1-q} \bar{a}_n^{(\alpha)} \right] + \delta_{mn} \left[\frac{q}{1-q} \bar{b}_n^{(\alpha)} \right]. \quad (5.4)$$

Here $\bar{a}_n^{(\alpha)} := a_n^{(\alpha)}(q\kappa)$, $\bar{b}_n^{(\alpha)} := b_n^{(\alpha)}(q\kappa)$.

From these results, we can exact useful information by specifying the value of m :

Case 1: $m < n - 1 \Leftrightarrow m + 2 \leq n$

We find, by induction, $\xi_{mn}^{(\alpha)}(\kappa) = 0$, if $m \leq n - 2$. Consequently, there are only two terms in the q -derivative (w.r.t κ variable) of the q -Laguerre orthonormal polynomials,

$$\mathcal{D}_q^\kappa p_n^{(\alpha)}(x, \kappa) = p_n^{(\alpha)}(x, \kappa) \xi_{nn}^{(\alpha)}(\kappa) + p_{n-1}^{(\alpha)} \xi_{n-1,n}^{(\alpha)}(\kappa). \quad (5.5)$$

Case 2: $m = n - 1$

In this case, we relate the two Fourier coefficients as follows:

$$\xi_{n-1,n}^{(\alpha)} = \frac{\frac{q}{1-q} \bar{a}_n^{(\alpha)}}{1 - \lambda \xi_{n-1,n-1}^{(\alpha)}} = \left(\frac{q}{1-q} \right) \frac{a_n^{(\alpha)}}{1 - \lambda \xi_{nn}^{(\alpha)}}. \quad (5.6)$$

Case 3: $m = n$

By suitable rearrangements, we derive a recursive equation relating the diagonal

Fourier coefficients $\xi_{nn}^{(\alpha)}$ to the recursive coefficients $a_n^{(\alpha)}, b_n^{(\alpha)}$ as follows:

$$(1 - \lambda \xi_{nn}^{(\alpha)})^2 + \frac{(q \kappa \bar{a}_n^{(\alpha)})^2}{(1 - \lambda \xi_{n-1,n-1}^{(\alpha)})^2} = 1 - q \kappa \bar{b}_n^{(\alpha)}. \quad (5.7)$$

By defining a sequence of functions in κ , which consist of alternating products/ratios of the recursive coefficients,

$$f_n^{(\alpha)}(\kappa) := \begin{cases} \left[\frac{\kappa \frac{a_1^{(\alpha)}(\kappa) a_3^{(\alpha)}(\kappa) \cdots a_n^{(\alpha)}(\kappa)}{a_2^{(\alpha)}(\kappa) a_4^{(\alpha)}(\kappa) \cdots a_{n-1}^{(\alpha)}(\kappa)} \right]^2, & \text{for odd } n \\ \left[\frac{a_2^{(\alpha)}(\kappa) a_4^{(\alpha)}(\kappa) \cdots a_n^{(\alpha)}(\kappa)}{a_1^{(\alpha)}(\kappa) a_3^{(\alpha)}(\kappa) \cdots a_{n-1}^{(\alpha)}(\kappa)} \right]^2, & \text{for even } n \end{cases}, \quad (5.8)$$

$$f_0^{(\alpha)}(\kappa) := 1. \quad (5.9)$$

One can check that

$$f_{n-1}^{(\alpha)}(\kappa) f_n^{(\alpha)}(\kappa) = [\kappa a_n^{(\alpha)}(\kappa)]^2.$$

If we repackage the recursive equation, Eq.(5.7), by introducing another variable

$$Y_n^{(\alpha)} := \frac{1}{\bar{f}_n^{(\alpha)}} (1 - \lambda \xi_{nn}^{(\alpha)})^2, \quad (5.10)$$

or equivalently,

$$\xi_{nn}^{(\alpha)} = \frac{1}{\lambda} \left(1 \pm \sqrt{\bar{f}_n^{(\alpha)} Y_n^{(\alpha)}} \right), \quad (5.11)$$

we have

$$Y_n^{(\alpha)} + \frac{1}{Y_{n-1}^{(\alpha)}} = \frac{1 - q\kappa \bar{b}_n^{(\alpha)}}{\bar{f}_n^{(\alpha)}}. \quad (5.12)$$

Hence, $Y_n^{(\alpha)}$ can be solved as a continued fraction in terms of the recursive coefficients $a_n^{(\alpha)}$ and $b_n^{(\alpha)}$.

Next, we study the q -difference equations describing the κ dependence of the recursive coefficients $a_n^{(\alpha)}(\kappa), b_n^{(\alpha)}(\kappa)$ associated with the generalized q -Laguerre ensemble. In the classical case, such equations correspond to the Lax pair formulation of the Toda equations. Hence, our results provide a q -generalization of the classical Toda equation.

To begin with, we compute the q -derivative with respect to the κ variable on the action of position operator, $x p_n^{(\alpha)}(x, \kappa)$, in two ways:

We first compute the q -derivative with respect to κ on the recursive relation,

$$\begin{aligned} \mathcal{D}_q^\kappa [x p_n^{(\alpha)}(x, \kappa)] &= \mathcal{D}_q^\kappa [a_{n+1}^{(\alpha)}(\kappa) p_{n+1}^{(\alpha)}(x, \kappa) + b_n^{(\alpha)}(\kappa) p_n^{(\alpha)}(x, \kappa) + a_n^{(\alpha)}(\kappa) p_{n-1}^{(\alpha)}(x, \kappa)] \\ &= [\mathcal{D}_q^\kappa a_{n+1}^{(\alpha)} + \xi_{n+1, n+1}^{(\alpha)} \bar{a}_{n+1}^{(\alpha)}] p_{n+1}^{(\alpha)} + [\mathcal{D}_q^\kappa b_n^{(\alpha)} + \xi_{n, n+1}^{(\alpha)} \bar{a}_{n+1}^{(\alpha)} + \xi_{nn}^{(\alpha)} \bar{b}_n^{(\alpha)}] p_n^{(\alpha)} \\ &+ [\mathcal{D}_q^\kappa a_n^{(\alpha)} + \xi_{n-1, n-1}^{(\alpha)} \bar{a}_n^{(\alpha)} + \xi_{n-1, n}^{(\alpha)} \bar{b}_n^{(\alpha)}] p_{n-1}^{(\alpha)} + [\xi_{n-2, n-1}^{(\alpha)} \bar{a}_n^{(\alpha)}] p_{n-2}^{(\alpha)}. \end{aligned} \quad (5.13)$$

Here $\xi_{mn}^{(\alpha)}$ are the Fourier coefficients(matrix elements) of Eq. (5.1),

$$\bar{a}_n^{(\alpha)} := a_n^{(\alpha)}(q\kappa), \bar{b}_n^{(\alpha)} := b_n^{(\alpha)}(q\kappa)$$

are the rescaled recursive coefficients, and we suppress the dependence on x, κ for simplicity.

On the other hand, since $\mathcal{D}_q^\kappa$ commutes with the position operator x , we first compute the q -derivative(w.r.t κ variable) of the orthonormal polynomials and then apply the position operator.

$$\begin{aligned} x \mathcal{D}_q^\kappa [p_n^{(\alpha)}(x, \kappa)] &= [\xi_{nn}^{(\alpha)} a_{n+1}^{(\alpha)}] p_{n+1}^{(\alpha)} + [\xi_{n-1, n}^{(\alpha)} a_n^{(\alpha)} + \xi_{nn}^{(\alpha)} b_n^{(\alpha)}] p_n^{(\alpha)} \\ &+ [\xi_{nn}^{(\alpha)} a_n^{(\alpha)} + \xi_{n-1, n}^{(\alpha)} b_{n-1}^{(\alpha)}] p_{n-1}^{(\alpha)} + [\xi_{n-1, n}^{(\alpha)} a_{n-1}^{(\alpha)}] p_{n-2}^{(\alpha)}. \end{aligned} \quad (5.14)$$

By comparing the corresponding coefficients of each orthonormal polynomials as calculated in Eqs.(5.13), (5.14), we get the following set of relations:

$$\mathcal{D}_q^\kappa a_n^{(\alpha)} = \xi_{n-1, n-1}^{(\alpha)} a_n^{(\alpha)} - \xi_{nn}^{(\alpha)} \bar{a}_n^{(\alpha)}, \quad (5.15)$$

$$\mathcal{D}_q^\kappa b_n^{(\alpha)} = \xi_{nn}^{(\alpha)} (b_n^{(\alpha)} - \bar{b}_n^{(\alpha)}) + [\xi_{n-1, n}^{(\alpha)} a_n^{(\alpha)} - \xi_{n, n+1}^{(\alpha)} \bar{a}_{n+1}^{(\alpha)}], \quad (5.16)$$

$$\mathcal{D}_q^\kappa a_n^{(\alpha)} = \xi_{n-1, n}^{(\alpha)} (b_{n-1}^{(\alpha)} - \bar{b}_n^{(\alpha)}) + [\xi_{nn}^{(\alpha)} a_n^{(\alpha)} - \xi_{n-1, n-1}^{(\alpha)} \bar{a}_n^{(\alpha)}], \quad (5.17)$$

$$\xi_{n-2, n-1}^{(\alpha)} \bar{a}_n^{(\alpha)} = \xi_{n-1, n}^{(\alpha)} a_{n-1}^{(\alpha)}. \quad (5.18)$$

Note that the last result allows us to replace the rescaled recursive coefficients in terms of the Fourier coefficients and the unscaled recursive coefficients

$$a_n^{(\alpha)}(q\kappa) = \frac{\xi_{n-1,n}^{(\alpha)}(\kappa)}{\xi_{n-2,n-1}^{(\alpha)}(\kappa)} a_{n-1}^{(\alpha)}(\kappa) = \frac{1 - \lambda \xi_{n-1,n-1}^{(\alpha)}}{1 - \lambda \xi_{nn}^{(\alpha)}} a_n^{(\alpha)}(\kappa). \quad (5.19)$$

We have also checked that Eqs.(5.15), (5.17) are compatible. Finally, after some manipulations, we get the coupled q -difference equations.

$$\mathcal{D}_q^\kappa a_n^{(\alpha)}(\kappa) = \frac{\xi_{n-1,n-1}^{(\alpha)}(\kappa) - \xi_{nn}^{(\alpha)}(\kappa)}{1 - \lambda \xi_{nn}^{(\alpha)}(\kappa)} a_n^{(\alpha)}(\kappa) \quad (5.20)$$

$$\mathcal{D}_q^\kappa b_n^{(\alpha)}(\kappa) = \frac{q}{1-q} \left[\left(\frac{a_n^{(\alpha)}(\kappa)}{1 - \lambda \xi_{nn}^{(\alpha)}} \right)^2 - \left(\frac{a_{n+1}^{(\alpha)}(\kappa)}{1 - \lambda \xi_{n+1,n+1}^{(\alpha)}} \right)^2 \right]. \quad (5.21)$$

5.2. q -Difference equations for the recursive coefficients of the q -Hermite orthonormal polynomials.

In this section, we shall derive the q -difference equation for the recursive coefficients of the q -Hermite orthonormal polynomials. The starting point of our derivation is to compute the Fourier coefficients of the q -derivative of the q -Hermite orthonormal polynomials with respect to parameter κ ,

$$\mathcal{D}_q^\kappa P_n^{(\alpha)}(x; \kappa) = \sum_{j=0}^n P_j^{(\alpha)}(x; \kappa) \Xi_{jn}^{(\alpha)}(\kappa), \quad (n-j \text{ is even}). \quad (5.22)$$

Note that, by recalling the definition of the q -derivative, (A.2), we can also transform the equation above into the Fourier expansion of the q -evolved q -Hermite orthonormal polynomials with respect to parameter κ ,

$$P_n^{(\alpha)}(x; q\kappa) = [1 - \lambda \Xi_{nn}^{(\alpha)}] P_n^{(\alpha)}(x; \kappa) - \lambda \sum_{j=0}^{n-1} P_j^{(\alpha)}(x; \kappa) \Xi_{jn}^{(\alpha)}(\kappa). \quad (5.23)$$

Our first task is to derive a set of algebraic equations for the Fourier coefficients $\Xi_{jn}^{(\alpha)}(\kappa)$. By taking the q -derivative (w.r.t κ) on the orthonormal condition

$$\mathcal{D}_q^\kappa \left(\int_{-1}^1 P_m^{(\alpha)}(x; \kappa) P_n^{(\alpha)}(x; \kappa) \omega^{(\alpha)}(x; \kappa) d_q x \right) = 0, \quad (5.24)$$

we get (assuming $m \leq n$)

$$\begin{aligned}
& \int_{-1}^1 (\mathcal{D}_q^\kappa P_m^{(\alpha)}(x; \kappa)) P_n^{(\alpha)}(x; \kappa) \omega^{(\alpha)}(x; \kappa) d_q x \\
& + \int_{-1}^1 P_m^{(\alpha)}(x; q\kappa) (\mathcal{D}_q^\kappa P_n^{(\alpha)}(x; \kappa)) \omega^{(\alpha)}(x; \kappa) d_q x \\
& + \int_{-1}^1 P_m^{(\alpha)}(x; q\kappa) P_n^{(\alpha)}(x; q\kappa) (\mathcal{D}_q^\kappa \omega^{(\alpha)}(x; \kappa)) d_q x = 0.
\end{aligned} \tag{5.25}$$

Substituting the Fourier expansions, Eqs.(5.22) and (5.23), into the first two terms of (5.25), and using the Pearson relation (in the κ variable) for the q -Hermite weight, we get

$$\begin{aligned}
& \delta_{mn} \Xi_{nn}^{(\alpha)} + (1 - \lambda \Xi_{mn}^{(\alpha)}) \Xi_{mn}^{(\alpha)} - \lambda \sum_{j=0}^{m-1} \Xi_{jm}^{(\alpha)} \Xi_{jn}^{(\alpha)} \\
& + \delta_{mn} \left(\frac{q^2 \kappa}{q-1} \right) \left\{ \left(\bar{A}_{n+1}^{(\alpha)} \right)^2 + \left(\bar{A}_n^{(\alpha)} \right)^2 \right\} + \delta_{m,n-2} \left(\frac{q^2 \kappa}{q-1} \right) \bar{A}_{n-1}^{(\alpha)} \bar{A}_n^{(\alpha)} = 0.
\end{aligned} \tag{5.26}$$

In order to illustrate the content of this equation, we consider the following specializations:

Case 1: $m < n - 2 \Leftrightarrow m + 3 \leq n$

In this case, the master equation reduces to

$$(1 - \lambda \Xi_{mm}^{(\alpha)}) \Xi_{mn}^{(\alpha)} - \lambda \sum_{j=0}^{m-1} \Xi_{jm}^{(\alpha)} \Xi_{jn}^{(\alpha)} = 0.$$

By the mathematical induction, we show that $\Xi_{mn}^{(\alpha)} = 0$, if $3 \leq n - m$. Consequently, the Fourier expansion of the q -derivative (w.r.t κ variable) on the q -Hermite orthonormal polynomials only consist of two terms:

$$\mathcal{D}_q^\kappa P_n^{(\alpha)}(x, \kappa) = P_n^{(\alpha)}(x, \kappa) \Xi_{nn}^{(\alpha)}(\kappa) + P_{n-2}^{(\alpha)}(x, \kappa) \Xi_{n-2,n}^{(\alpha)}(\kappa). \tag{5.27}$$

Case 2: $m = n - 2 \Leftrightarrow m - 1 = n - 3$

In this case, the master equation reduces to

$$(1 - \lambda \Xi_{n-2,n-2}^{(\alpha)}) \Xi_{n-2,n}^{(\alpha)} - \lambda \sum_{j=0}^{n-3} \Xi_{j,n-2}^{(\alpha)} \Xi_{jn}^{(\alpha)} + \frac{q^2 \kappa}{q-1} \bar{A}_n^{(\alpha)} \bar{A}_{n-1}^{(\alpha)} = 0.$$

Since we have showed that $\Xi_{jn}^{(\alpha)} = 0$ for $j \leq n - 3$ in Case 1, we can use the equation above to express the off-diagonal Fourier coefficient in terms of the diagonal ones:

$$\Xi_{n-1,n+1}^{(\alpha)} = \left(\frac{q^2 \kappa}{1-q} \right) \frac{\bar{A}_{n+1}^{(\alpha)} \bar{A}_n^{(\alpha)}}{1 - \lambda \Xi_{n-1,n-1}^{(\alpha)}}. \tag{5.28}$$

Case 3: $m = n - 1, m - 1 = n - 2$

Due to the parity preserving property associated with the q -Hermite ensemble,

$\Xi_{n-1,n}^{(\alpha)} = 0$, we have no constraint in this case.

Case 4: $m = n$

In this case, we have

$$\Xi_{nn}^{(\alpha)} + (1 - \lambda \Xi_{nn}^{(\alpha)}) \Xi_{nn}^{(\alpha)} - \lambda (\Xi_{n-2,n}^{(\alpha)})^2 = \left(\frac{q^2 \kappa}{1 - q} \right) \left[(\bar{A}_{n+1}^{(\alpha)})^2 + (\bar{A}_n^{(\alpha)})^2 \right]. \quad (5.29)$$

After suitable rearrangement, we get

$$(1 - \lambda \Xi_{nn}^{(\alpha)})^2 + \frac{(q\kappa)^4 (\bar{A}_n^{(\alpha)})^2 (\bar{A}_{n-1}^{(\alpha)})^2}{(1 - \lambda \Xi_{n-2,n-2}^{(\alpha)})^2} = 1 - (q\kappa)^2 \left\{ (\bar{A}_{n+1}^{(\alpha)})^2 + (\bar{A}_n^{(\alpha)})^2 \right\}. \quad (5.30)$$

Similar to the case of the q -Laguerre ensemble Eq.(5.7), we can solve $(1 - \lambda \Xi_{nn}^{(\alpha)})^2$ as a continued fraction in terms of the recursive coefficient $\bar{A}_n^{(\alpha)}$.

To summarize, the previous calculations show that we can compute the Fourier coefficients of $\mathcal{D}_q^\kappa P_n^{(\alpha)}(x; \kappa)$ in terms of the recursive coefficients, $A_n^{(\alpha)}(\kappa)$, of the q -Hermite orthonormal polynomial. Our next step is to deduce the q -difference equation for these recursive coefficients. Similar to the q -Laguerre case, Eq.(5.13), we can compute the $\mathcal{D}_q^\kappa$ derivative on the product $x P_n^{(\alpha)}(x; \kappa)$ in two ways.

$$\begin{aligned} \mathcal{D}_q^\kappa [x P_n^{(\alpha)}(x; \kappa)] &= \mathcal{D}_q^\kappa [A_{n+1}^{(\alpha)}(\kappa) P_{n+1}^{(\alpha)}(x; \kappa) + A_n^{(\alpha)}(\kappa) P_{n-1}^{(\alpha)}(x; \kappa)] \\ &= x \left[\sum_{j=0}^n P_j^{(\alpha)}(x; \kappa) \Xi_{jn}^{(\alpha)}(\kappa) \right] \\ &= \sum_{j=0}^n [A_{j+1}^{(\alpha)}(\kappa) P_{j+1}^{(\alpha)}(x; \kappa) + A_j^{(\alpha)}(\kappa) P_{j-1}^{(\alpha)}(x; \kappa)] \Xi_{jn}^{(\alpha)}(\kappa). \end{aligned} \quad (5.31)$$

By comparing the coefficients of $P_{n+1}^{(\alpha)}(x; \kappa)$ of the first and the third lines of the previous equation, we get the following results:

$$\mathcal{D}_q^\kappa A_{n+1}^{(\alpha)}(\kappa) = \Xi_{nn}^{(\alpha)}(\kappa) A_{n+1}^{(\alpha)}(\kappa) - \Xi_{n+1,n+1}^{(\alpha)}(\kappa) A_{n+1}^{(\alpha)}(q\kappa), \quad (5.32)$$

which implies

$$\mathcal{D}_q^\kappa A_n^{(\alpha)}(\kappa) = \frac{\Xi_{n-1,n-1}^{(\alpha)}(\kappa) - \Xi_{nn}^{(\alpha)}(\kappa)}{1 + \kappa(q-1)\Xi_{nn}^{(\alpha)}(\kappa)} A_n^{(\alpha)}(\kappa). \quad (5.33)$$

6. SUMMARY AND CONCLUSION

In this paper, we study fundamental properties of the generalized q -Laguerre/Hermite ensembles with a deformation parameter κ .

For the special value $\kappa = 1$, we give explicit solutions of their recursive coefficients for the orthonormal polynomial systems. In addition, to show that these ensembles provide "good" basis for a realization of the Heisenberg algebra, we also calculate the matrix elements (Fourier coefficients) of the dilation operator with respect to orthonormal polynomial basis explicitly.

In view of the intimate connections between matrix models and orthogonal polynomial systems [8], we compute the Hankel determinants as products of the recursive coefficients explicitly. The physical consequences for identifying the Hankel determinants as partition functions of these matrix models with non-polynomial type potential may worth further explorations.

Finally, we examine the deformation of the q -Laguerre/Hermite systems in the form of q -difference equations dictating the κ dependence of the recurrence coefficients and Fourier coefficients. This provides a q -generalization of the Toda equations in the Lax pair formulation [10].

ACKNOWLEDGEMENT

This research project was initiated in a summer visit to Institute of Mathematics at Academia Sinica in 2015. Both authors would like to thank Derchyi Wu and Chueh-Hsin Chang, and Mourad E. H. Ismail for instructive discussions. C.T. would like to thank Derchyi Wu for invitation and hospitality. The research work of C.T. is partially supported by the grant from Academia Sinica for summer visit, and in parts supported by the MOST research grant 104-2112-M-029-001. The research work of H.F. is partially supported by the MOST research grant 104-2115-M-001-001-MY2.

APPENDIX A. SOME BASIC DEFINITIONS AND RELATIONS FOR q -ANALYSIS ($0 < q < 1$)

In this section, we collect some basic definitions and formulas which are relevant to our discussions.

The q -integral for a function $f(x)$ over the region $x \in [0, a]$ is defined as

$$F(a) := \int_0^a f(x) d_q x := a(1-q) \sum_{k=0}^{\infty} f(aq^k) q^k. \quad (\text{A.1})$$

This is compatible with the definition of the q -derivative

$$\mathcal{D}_q f(x) := \frac{f(qx) - f(x)}{qx - x} = \frac{f(x) - f(qx)}{x(1-q)} \quad (\text{A.2})$$

in the following senses:

(1) Fundamental theorem of the q -calculus

$$\mathcal{D}_q F(a) = f(a), \quad (\text{A.3})$$

$$\int_0^a [\mathcal{D}_q f(x)] d_q x = f(a) - f(0). \quad (\text{A.4})$$

(2) The linear change of variables can be implemented in q -integral:

$$\int_0^a f(cx) d_q x = \frac{1}{c} \int_0^{ca} f(y) d_q y. \quad (\text{A.5})$$

There are some subtleties associated with the q -derivative, in particular, the q -Leibnitz rule is given as

$$\begin{aligned} \mathcal{D}_q [f(x)g(x)] &= \frac{f(qx)g(qx) - f(x)g(x)}{(q-1)x} \\ &= f(qx)[\mathcal{D}_q g(x)] + [\mathcal{D}_q f(x)]g(x) \\ &= [\mathcal{D}_q f(x)]g(qx) + f(x)[\mathcal{D}_q g(x)]. \end{aligned} \quad (\text{A.6})$$

APPENDIX B. ON THE q -DEFORMATION OF THE HEISENBERG ALGEBRA

In the classical case, the Heisenberg algebra is generated by two operators: $\frac{d}{dx}$ (translation) and x (position). They satisfy the commutation relation,

$$\left[\frac{d}{dx}, x \right] = 1. \quad (\text{B.1})$$

We can also include the dilation operator $\mathcal{S} := x \frac{d}{dx}$, such that

$$[\mathcal{S}, x] = \left[x \frac{d}{dx}, x \right] = x, \quad (\text{B.2})$$

and

$$\left[\mathcal{S}, \frac{d}{dx} \right] = \left[x \frac{d}{dx}, \frac{d}{dx} \right] = -\frac{d}{dx}. \quad (\text{B.3})$$

In this way, we can view the position (x) and translation ($\frac{d}{dx}$) operators as raising and lowering operators with respect to the eigenfunctions of dilation.

To study the q -deformation of the Heisenberg algebra, which is useful in our computations of the recursive coefficients for the q -Laguerre and q -Hermite ensembles, we compute the action of three commutators on a general function $f(x)$: ($\mathcal{S}_q := x\mathcal{D}_q$)

Theorem B.1. *The q -difference realization of the q -Heisenberg algebra is given by*

$$[\mathcal{D}_q, x]f(x) = f(qx)$$

$$[\mathcal{S}_q, x]f(x) = xf(qx)$$

$$[\mathcal{S}_q, \mathcal{D}_q]f(x) = -\mathcal{D}_q f(qx).$$

Proof.

$$\begin{aligned} [\mathcal{D}_q, x]f(x) &= \mathcal{D}_q(xf(x)) - x(\mathcal{D}_q f(x)) \\ &= \frac{qx f(qx) - xf(x)}{(q-1)x} - \frac{f(qx) - f(x)}{q-1} \\ &= f(qx). \end{aligned}$$

$$\begin{aligned} [\mathcal{S}_q, x]f(x) &= \mathcal{S}_q(xf(x)) - x(\mathcal{S}_q f(x)) \\ &= \frac{qx f(qx) - xf(x)}{q-1} - x \frac{f(qx) - f(x)}{q-1} \\ &= xf(qx). \end{aligned} \quad (\text{B.4})$$

$$\begin{aligned}
[\mathcal{S}_q, \mathcal{D}_q]f(x) &= \mathcal{S}_q(\mathcal{D}_q f(x)) - \mathcal{D}_q(\mathcal{S}_q f(x)) \\
&= \mathcal{S}_q\left(\frac{f(qx) - f(x)}{(q-1)x}\right) - \mathcal{D}_q\left(\frac{f(qx) - f(x)}{q-1}\right) \\
&= -\frac{f(q^2x)}{q(q-1)x} + \frac{f(qx)}{q(q-1)x} \\
&= -\mathcal{D}_q f(qx).
\end{aligned} \tag{B.5}$$

□

Theorem B.2. *For any q -weight function, the diagonal matrix elements (or the Fourier coefficients) of the q -dilation operator in terms of orthonormal polynomial basis, defined by*

$$\mathcal{S}_q p_n(x) = \sum_{j=0}^n p_j(x) c_{jn},$$

have an universal expression:

$$c_{nn} = \frac{1 - q^n}{1 - q}.$$

APPENDIX C. A SHORT NOTE FOR THE q -EXPONENTIAL FUNCTION

The q -deformed exponential function is defined either by an infinite product or as a power series in x ,

$$e_q(x) := \frac{1}{\prod_{k=0}^{\infty} (1 - q^k x)} \quad (\text{C.1})$$

$$= 1 + \sum_{n=1}^{\infty} \frac{x^n}{\prod_{k=1}^n (1 - q^k)} \quad (\text{C.2})$$

$$= 1 + \sum_{n=1}^{\infty} \frac{x^n}{(1 - q)(1 - q^2) \cdots (1 - q^n)}. \quad (\text{C.3})$$

It is easy to see that, using the power series form, Eq. (C.3), as $\epsilon := 1 - q \rightarrow 0$, we can connect this q -exponential function to the classical exponential and gaussian functions.

$$\lim_{\epsilon \rightarrow 0} e_{1-\epsilon}(\epsilon x) = \lim_{\epsilon \rightarrow 0} \left(1 + \sum_{n=1}^{\infty} \frac{\epsilon^n x^n}{\epsilon \cdot 2\epsilon \cdots n\epsilon} \right) = e^x.$$

$$\lim_{q \rightarrow 1} e_{q^2}((1 - q^2)x^2) = \lim_{\epsilon \rightarrow 0} \left(1 + \sum_{n=1}^{\infty} \frac{(2\epsilon)^n x^{2n}}{2\epsilon \cdot 4\epsilon \cdots 2n\epsilon} \right) = e^{x^2}. \quad (\text{C.4})$$

REFERENCES

- [1] Ismail, Mourad E. H.; Stanton, Dennis; Viennot, Gérard, *The combinatorics of q -Hermite polynomials and the Askey-Wilson integral*, European J. Combin. 8 (1987), no. 4, 379-392.
- [2] Ismail, Mourad E. H., *Orthogonal Polynomials, Their Recursions and Functional Equations*, Symmetries and Integrability of Difference Equations, Cambridge Univ. Press, London Math. Soc. Lecture Note Series: 381, 2011.
- [3] Clarkson, Peter A., *Recurrence coefficients for discrete orthonormal polynomials and the Painlevé equations*, J. Phys. A 46 (2013), no. 18, 185205, 18 pp.
- [4] Boelen, Lies; Van Assche, Walter, *Discrete Painlevé equations for recurrence coefficients of semiclassical Laguerre polynomials*, Proc. Amer. Math. Soc. 138 (2010), no. 4, 1317-1331
- [5] Van Assche, Walter *Discrete Painlevé equations for recurrence coefficients of orthogonal polynomials*, Difference equations, special functions and orthogonal polynomials, 687-725, World Soc. Publ., Hackensack, NJ, 2007.
- [6] Kasraoui, Anisse; Stanton, Dennis; Zeng, Jiang, *The combinatorics of Al-Salam-Chihara q -Laguerre polynomials*, Adv. in Appl. Math. 47 (2011), no. 2, 216-239.
- [7] Simion, R.; Stanton, D., *Specializations of generalized Laguerre polynomials*, SIAM J. Math. Anal. 25 (1994), no. 2, 712-719.
- [8] P. Di Francesco; P. H. Ginsparg; J. Zinn-Justin, *2-D Gravity and random matrices*, Phys. Rept. **254**, 1 (1995) [hep-th/9306153].
- [9] K. Fujikawa, *Path integral of the hydrogen atom, Jacobi's principle of least action and one-dimensional quantum gravity*, Nucl. Phys. B **484**, 495 (1997) [hep-th/9602080].
- [10] H. Flaschka, *The Toda lattice. II. Existence of integrals*, Phys. Rev. B 9, p.1924, 1974